arXiv:astro-ph/0010315v1 16 Oct 2000

Probing the Substellar Regime with SIRTF

Eduardo L. Martín, Wolfgang Brandner, David Jewitt, Theodore Simon, and Richard Wainscoat
Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822
`ege@ifa.hawaii.edu,brandner@ifa.hawaii.edu,jewitt@ifa.hawaii.edu,simon@ifa.hawaii.edu,`
`rjw@ifa.hawaii.edu`

Mark Marley
Space Physics Research Institute, NASA Ames Research Center, Moffett Field, CA 94035
`mmarley@mail.arc.nasa.gov`
and

Christopher Gelino
Department of Astronomy, New Mexico State University, Las Cruces, NM 880033
`crom@nmsu.edu`

ABSTRACT

One of the main scientific drivers of the Space InfraRed Telescope Facility (SIRTF) is the search for brown dwarfs and extrasolar superplanets. We discuss observational strategies for identification of these objects, and conclude that an optimal strategy is a wide IRAC survey (18 deg^2) with a 5 σ sensitivity of 3.9 μJy in channel 2 (M$\sim$19.1^m). For this sensitivity, we provide estimates of the number of low mass brown dwarfs and isolated planets detected per square degree for power-law mass functions with α=1.5, and 1.0. Shallower surveys covering a larger area are inefficient because of large overheads and detector noise. Deeper surveys covering a smaller area become more and more affected by crowding with galaxies. A survey like the one that we propose would determine the field mass function down to a few Jupiter masses through the identification of a large sample of brown dwarfs and isolated planets. The proposed SIRTF survey would also allow the first detection of ultracool substellar objects with temperatures between 700 K and 200 K. The cooling curves of substellar objects with masses less than 20 Jupiters imply that they should spend most of their lifetimes at temperatures below 700 K. Preliminary models indicate that their atmospheres could be dominated by water clouds, which would diminish their optical and near-infrared fluxes. The properties of those objects are still completely unexplored.

Subject headings: techniques:photometric – solar neighborhood – Galaxy: structure – infrared: general – stars: low mass, brown dwarfs – planetary systems

1. Introduction

The past five years have witnessed the remarkable birth of a new research field of astronomy — the physical study of substellar objects outside the solar system. Large scale optical and near-infrared surveys of the sky (DENIS, 2MASS and SDSS) have revealed very cool objects that are close to us but nevertheless were unknown. The overwhelming majority of substellar objects in the field, however, cannot be detected by ground-based surveys;

such objects cool very rapidly with time and become extremely faint at wavelengths shorter than about 4 μm. SIRTF (Fanson et al. 1998) will allow to probe deeper in the substellar regime, extending the census of substellar objects in the field to lower masses and temperatures (see Figure 1).

The separation between brown dwarfs and planets is a matter of convention. We concur with other authors (e.g., Basri & Marcy 1997; Oppenheimer et al. 2000) in arguing that a definition based on the well known laws of gravitation and nuclear physics is more reliable than a definition based on the origin of the objects. Thus, we adopt the following definitions: A brown dwarf is an object unable to stabilize on the hydrogen burning main-sequence, but able to fuse deuterium into helium. For solar metallicity the mass range of brown dwarfs is between 0.075 $M_\odot$ and 0.013 $M_\odot$ (Baraffe et al. 1998, Saumon et al. 1996). Planets are objects unable to fuse deuterium, i.e., they have masses lower than 0.013 $M_\odot$. "Isolated planets" are planets that are not bound to any particular star. Recent results suggest that this kind of planets may be numerous in the disk of the Galaxy (Béjar et al. 2000; Zapatero Osorio et al. 2000). The smooth transition of the mass function from the brown dwarfs to the isolated planets indicates that they have a common origin, which is much like that of star formation from fragmentation and collapse of molecular clouds, but not like the planets in our solar system (accretion in a protoplanetary disk). Despite of their different origins, isolated planets may have the same basic properties (overall chemical composition, luminosity, temperature) as their counterparts orbiting stars.

Studies show that the number of substellar objects increases towards lower masses. The functional form of this relationship, the Initial Mass Function (IMF hereafter), can be approximated by a power law $dN/dM \sim M^{-\alpha}$ (Salpeter 1955). For masses between 0.1 $M_\odot$ and 0.01 $M_\odot$ (i.e., $\sim$ 100 and 10 M_{Jup}, respectively), several surveys in young open clusters have found $\alpha = 1.0 \pm 0.5$. (Béjar et al. 2000, Hillenbrand & Carr 2000; Martín et al. 2000; Najita et al. 2000). The density of field brown dwarfs discovered by DENIS and 2MASS is also consistent with this mass function (Reid et al. 1999). The number of substellar objects in each cluster, or in the field, is still small. Hence the statistical uncertainties are large and it is not possible to conclude if there are significant variations in the IMF from one region to another. The cooling curves of brown dwarfs and planets imply that they spend a large part of their lives at extremely low luminosities and cool temperatures. The launch of the SIRTF satellite provides an unprecedented opportunity for detecting ultracool substellar objects.

This paper is organized as follows: Section 2 gives a brief description of the expected spectra of ultracool substellar objects and justifies the need for SIRTF to detect them; Section 3 discusses observational strategies to detect substellar objects with SIRTF; Section 4 provides estimates of the counts of substellar objects in the disk that can be detected with SIRTF; and Section 5 summarizes the main conclusions.

2. The need for SIRTF: Infrared Spectra of Ultracool Substellar Objects

We have used calculations for solar composition substellar objects. These models will be described in detail in a forthcoming paper (Marley et al. 2001, in preparation). We limit this presentation to a brief discussion of results that are relevant for our purposes.

With decreasing temperature the molecular bands of methane and water depress the flux at optical and near-infrared wavelengths. At temperatures below $\sim$700 K, substellar objects emit very little flux shortward of 4 μm (see Figure 2). In fact, the coolest known brown dwarfs, namely Gl 570 D, has a temperature of 790±50 K (Burgasser et al. 2000). Its spectrum is similar to that of Gl 229 B.

It is unlikely that the DENIS and 2MASS surveys will detect any objects significantly cooler than Gl 570 D. With $J = 15\overset{\rm m}{.}33$ and $M_J = 16\overset{\rm m}{.}5$, this object is near the detection limit of 2MASS ($J \sim 16^m$; and, $J \sim 15^m$ for DENIS). A substellar object 200 K cooler would have an absolute J-band magnitude about 1.5 mag fainter (Burrows et al. 1997), and to be detected by these surveys it would have to be 2 times closer, i.e., at 3 pc. There are 6 stellar systems within 3 pc of the Sun, and there could be a similar number of brown dwarfs. However, the chances that 2MASS will detect them may be negligible if the near-infrared spectrum is depleted by strong water bands.

Preliminary models (Marley 2000), illustrated here in Figure 3, provide a glimpse of what these substellar objects may look like. Water clouds will first appear in objects with $T_{\rm eff}$ below $\sim$600 K. The clouds will first form as tenuous stratus clouds high in the atmosphere and then form lower in the atmosphere in cooler objects where they will be thicker and more substantial. For $T_{\rm eff}$ below $\sim$300 K, ammonia clouds will form. Jupiter's water clouds are hidden below two higher cloud layers and are poorly understood. Thus SIRTF provides the opportunity to detect the first substellar mass objects outside our solar system in which water clouds dominate the atmosphere.

Water clouds will substantially alter the infrared spectra of these objects. T-dwarfs like Gliese 229 B are blue in J-K. Once water clouds form, the J-K color will move redward towards J-K 0 to 1 (Marley and Ackerman 2000). A new spectral class, P for example, could be necessary for classifying these objects cooler than the T dwarfs. Based on near-IR colors (JHK) alone, it will be very difficult to distinguish the P-type brown dwarfs from background M-stars with similar colors. Brown dwarfs with clear atmospheres, like the T-dwarfs, are bright at 4 to 5 um (Marley et al. 1996; Burrows et al. 1997). As with Jupiter, Marley and Ackerman (2000) find that the coolest brown dwarfs are also bright in this spectral region, even when water clouds are present (Figure 3). Thus a search with SIRTF at 4-5um has the potential of easily discovering such objects.

As illustrated in Figure 2, the IRAC (Fazio et al. 1998) passband at 4.5 μm (channel 2) offers the best sensitivity for detection of a 10 M_{Jup} object, particularly for old ages. 2MASS can detect a 10 M_{Jup} isolated planet only if it is about 0.1 Gyr old and is located at a distance of less than 10 pc. It is unlikely that such an object exists. On the other hand, SIRTF can provide a 5 σ detection in 720 s of a 10 M_{Jup} isolated planet with an age of 0.1 Gyr out to 350 pc at 4.5 μm and 200 pc at 3.6 μm. SIRTF can also detect a 10 M_{Jup} isolated planet at an age 1 Gyr up to a distance of 120 pc at 4.5 μm and 40 pc at 3.6 μm.

3. Observational Strategies for Detecting Substellar Objects with SIRTF

The DENIS, 2MASS and SDSS surveys have demonstrated that free-floating brown dwarfs outnumber brown dwarf companions to stars by about a factor of 10. These surveys combined have already discovered more than 200 free-floating brown dwarfs (Burgasser et al. 1999; Delfosse et al. 1999; Fan et al. 2000; Gizis et al. 2000; Kirkpatrick et al. 2000; Leggett et al. 2000; Martín et al. 1999), but only three companion brown dwarfs to stars (Burgasser et al. 2000; Kirkpatrick et al. 2000). Moreover, high-precision radial velocity surveys have revealed a brown dwarf desert within a few AU of G and K-type main-sequence stars (Halbwachs et al. 2000). They determine that there exists a secondary mass cutoff between 0.01 and 0.1 $M_\odot$. Other searches for brown dwarf companions to stars have found only a few outstanding examples in samples of hundreds of stars (Becklin & Zuckerman 1988; Oppenheimer 1999). These companions are very interesting but seem to be rare.

The spatial resolution of SIRTF (FWHM=2".3 with IRAC), and the lack of a coronograph, will not allow a Jupiter mass planet at 5 AU separation to be resolved, even for the closest stars. Jupiter-mass planets may be common around stars, but they are unlikely to be found outside a 20 AU radius of the parent star. The ice condensation region is located at about 5 AU, with little dependence on the spectral type of the star, and gravitational instabilities in the disk are unlikely to be effective beyond 20 AU (Boss 2000). Direct images of massive disks around T Tauri stars indicate that in general the density is too low to allow for the formation of a giant planet beyond 20 A.U. (e.g., Krist et al. 2000). On the other hand, the deep surveys of Orion and IC 348 suggest that free-floating planets probably are very numerous (Najita et al 2000; Zapatero Osorio et al. 2000).

These considerations lead us to propose that a wide field imaging survey with SIRTF is the best strategy to detect a large sample of substellar objects. We have used the SIRTF planning observations tool (SPOT) to simulate future observations. With total exposure times of 720 s per pointing (divided into 24 exposures of 30 s for dithering and cosmic ray removal), we could make an IRAC map covering 1 square degree in 33.3 hours of observation time. The 5 σ sensitivity for point sources under conditions of low background are: 2.91 μJy in channel 1, 3.88 μJy in channel 2, 11.3 μJy in channel 3, and 13.2 μJy in channel 4. At these flux limits, the main contribution to source confusion comes from background galaxies We have estimated the number of galaxies and stars at the IRAC passbands using the predictions of Franceschini et al. (1991) and a modified version of the galaxy model of Wainscoat et al. (1992). the recent 7μm ISOCAM counts obtained by Taniguchi et al. (1997). Our calculations show that in IRAC channel 2 (our most sensitive passband for substellar objects) we expect to detect approximately 3,800 stars (for high galactic latitude) and 78,000 galaxies per square degree. If each galaxy is assumed to occupy 3×3 pixels (3.6×3.6 arcsec), then the galaxies will cover approximately 7% of the pixels. At fainter detection limits, the crowding from galaxies is correspondingly greater. Thus, more sensitive observations offer relatively diminishing returns for a much greater investment of observing time.

According to the fluxes predicted by our synthetic spectra, substellar objects can be identified unambiguously from IRAC photometry. In fact, it is enough with a 5 σ detection in channel 2 and no detection in the other channels. As shown by Figure 4, any object with log F(4.5μm)/F(3.6μm) > 0.3 can be considered as a good brown dwarf or planet candidate. Neither stars nor galaxies contaminate the substellar locus. The location of flared and flat disks in Figure 4 is based on results on the spectral index for T Tauri star disks by Kenyon & Hartmann (1995). The location of 30 Doradus has been derived from ISO SWS measurements by Sturm et al. (2000), and the flux values for L* galaxies at z=0.1 to 5 are based on simulations by Fazio et al. (1998).

Figure 4 also indicates that Population III[1] brown dwarfs with effective temperatures below 400 K are readily identified from their colors alone. Still warmer (and more massive) Population III brown dwarfs occupy a region in the color-color plot that coincides with redshifted L* galaxies and with stars having debris disks which are intermediate between Vega-type disks and the disks around classical and weak-line T Tauri stars. Additional information will therefore be needed to make a positive identification. Two epochs of IRAC observations would facilitate an identification of the most nearby Population III brown dwarfs, based on their high proper motion (and possibly their parallax). More distant candidates for massive Population III brown dwarfs will have to be followed up with instruments with higher instrinsic angular resolution, e.g., ground-based 8-m to 10-m class telescopes, the Next Generation Space Telescope (NGST), the Terrestrial Planet Finder (TPF), or the Stratospheric Observatory for Infrared Astronomy (SOFIA).

4. Modelling the Number Counts of Substellar Objects in the Disk

We have estimated the number counts of substellar objects in the disk of the Galaxy using the following ingredients: (1) Theoretical fluxes computed from synthetic spectra (see Figure 2). (2) Sensitivities for low-background from the IRAC manual. We have adopted a 5σ confidence level for our detection limit in each channel. (3) A solar neighborhood density of 0.057 stellar systems/pc^3 (Reid et al. 1999). (4) Power-law mass functions with α=1.5, 1.0 and 0.5 from 100 to 1 M_{Jup} . (5) A constant star formation rate over the lifetime of the Galactic disk (from 0 to 10 Gyr). (6) An exponential vertical distribution for distances beyond 50 pc, and a scale height of 325 pc, consistent with the counts of M dwarfs.

A Monte Carlo code provided the cumulative number of substellar objects. In Figure 5, we show the cumulative distributions of objects with masses in the range 25 to 1 M_{Jup} for two different slopes of the IMF power-law. For an IMF with α = 1.5, we predict that SIRTF could detect about 30 isolated planets per square degree. For α = 1.0, the number of planets decreases to about 4 per square degree.

We consider the hypothetical case that 600 hours of SIRTF time would be dedicated for an IRAC survey to search for substellar objects. Using SPOT we have estimated that an area of 18 square degrees could be covered with 5 σ sensitivity of 3.88 μJy in channel 2. The cumulative distribution of substellar objects that such a survey would find (for $\alpha = 1.5$) is given in Figure 6. We compare with another survey with which is less sensitive but covers a larger area (64 deg^2). Both surveys require the same amount of SIRTF observing time according to the SPOT program. The larger survey is less efficient due to larger overheads and the contribution of detector noise when the exposure times are shorter. More than 150 substellar objects with temperatures cooler than 600 K could be revealed in 18 square degrees. If the IMF is shallower the number of objects would naturally be less. For $\alpha = 1.0$ we predict about 20 objects cooler than 600 K, and for $\alpha = 0.5$ we predict only three of them. SIRTF observations can constrain the field mass function down to masses below the deuterium limit by counting the number of ultracool substellar objects.

---

[1] Population III stars in general refers to the first generation of stars, i.e. objects which formed at zero-metallicity (e.g. Cojazzi et al. 2000, and references therein). Similarly, Population III brown dwarfs refers to substellar objects, which formed at zero-metallicity.

5. Conclusions

We have shown that IRAC channel 2 is very efficient for detecting ultracool substellar objects. We present an optimized observational strategy for detection of brown dwarfs and isolated planets with temperatures below 700 K. Preliminary models indicate that the atmospheres of these objects are dominated by water clouds, which make the objects fainter at optical and near-infrared wavelengths than previously thought. The 5 μm window is an excellent region for detecting them because their atmospheres are more transparent. Our strategy consists of a SIRTF survey with a 5 σ sensitivity of 3.88 μJy in IRAC channel 2. We show that substellar objects can easily be identified in an IRAC color-color diagram. Any object with $\log F(4.5\mu m)/F(3.6\mu m) > 0.3$ can be considered as a good brown dwarf or planet candidate. Neither stars nor galaxies contaminate the substellar locus. With our detection limit, we predict a density of 30 isolated planets per square degree for an IMF with a slope $\alpha = 1.5$. In 600 hours of SIRTF time, an area of 18 square degrees would be covered. We predict that such a survey would find about 540 isolated planets, and about 160 substellar objects with temperatures cooler than 600 K. Such objects cannot be detected with the current generation of near-infrared all-sky surveys (DENIS and 2MASS). For shallower IMF slopes, the number of substellar objects diminishes. Thus, the proposed survey can determine the field substellar IMF down to very low masses.

We would like to thank the following people for useful comments and discussions: Xavier Delfosse, Thierry Forveille, John Gizis, Pat Henry, Klaus Hodapp, Joseph Hora, Jeff Kuhn, Alan Tokunaga, Motohide Tamura, and Maria Rosa Zapatero Osorio. We thank France Allard and Isabelle Baraffe for sending their theoretical models so that we could compare them with the models used in this paper.

REFERENCES

Ackerman, A. & Marley, M. S. 2000, in preparation

Baraffe, I., Chabrier, C., Allard, F., & Hauschildt, P. H. 1998, A&A, 337, 403

Basri, G. & Marcy, G. 1997, in Star Formation Near and Far, Holt & Mundy, eds, New York AIP Press, 393, 228

Becklin, E. E. & Zuckerman, B. 1988, Nature, 336, 656

Béjar, V. J. S. et al. 2000, ApJ, submitted

Boss, A. P. 2000, ApJ, 536, L101

Burgasser, A. J. et al. 1999, ApJ, 522, L65

Burgasser, A. J. et al. 2000, ApJ, 531, L57

Burrows, A. et al. 1997, ApJ, 491, 856

Cojazzi, P., Bressan, A., Lucchin, F., Pantano, O., & Chavez, M. 2000, MNRAS, 315, L51

Delfosse, X. et al. 1997, A&A, 327, 25

Delfosse, X. et al. 1999, A&A, 344, 897

Delfosse, X. et al. 2000, A&A, in press

Fan, X. et al. 2000, ApJ, 119, 928

Fanson, J. L. et al. 1998, Proc SPIE, 3356, 478

Fazio, G. G. et al. 1998, Proc SPIE 3354, 1024

Franceschini, A., Toffolatti, L., Mazzei, P., Danese, L., & de Zotti, G. 1991, A&AS, 89, 285

Gizis, J. E., et al. 2000, AJ, 120, 1085

Halbwachs, J. L., Arenou, F., Mayor, M., Udry, S., & Queloz, D. 2000, A&A, 355, 581

Hillenbrand, L. A. & Carpenter, J. M. 2000, ApJ, 540, 236

Kenyon, S. J. & Hartmann, L. 1995, ApJS, 101, 117

Kirkpatrick, D. et al. 2000, AJ, 120, 447

Krist, J. E., Stapelfeldt, K. R., Ménard, F., Padgett, D. L., & Burrows, C. J. 2000, ApJ, 538, 793

Leggett, S. et al. 2000, ApJ, 536, L35

Marley, M. S. et al. 1996, Science, 272, 1919

Marley, M. S. 2000, XXIV IAU Meeting, Manchester

Martín, E. L., Basri, G. & Brandner, W. 1999a, Science, 283, 1718

Martín, E. L. et al. 1999b, AJ, 1727, 2466

Martín, E. L. et al. 2000, ApJ, in press

Najita, J. R., Tiede, G. P. & Carr, J. S. 2000, ApJ, in press

Oppenheimer, B. R. 1999, PhD Thesis, Caltech

Oppenheimer, B. R., Kulkarni, S. R. & Stauffer, J. 2000, in Protostars & Planets IV, ed. Mannings, V. et al. (Univ. of Arizona: Tucson), p. 1313

Reid, N. et al. 1999, ApJ, 521, 613

Salpeter, E. 1955, ApJ, 121, 161

Saumon, D., Hubbard, W. B., Burrows, A., Guillot, T., Lunine, J. I., & Chabrier, G. 1996, ApJ, 460, 993

Sturm, E. et al. 2000, A&A, 358, 481

Taniguchi, Y. et al. 1997, A&A, 328, L9

Wainscoat, R. et al. 1992, ApJS, 83, 111

Zapatero Osorio, M. R. et al. 2000, Science, in press

This 2-column preprint was prepared with the AAS LaTeX macros v5.0.

Fig. 1.— Parameter space occupied by Brown Dwarfs and Planets. Objects were selected from an $\alpha = 1.0$ power law distribution in mass with $0.01 < \text{age} < 10$ Gyr. Ground-based surveys are sensitive only to the minority of objects with $T_{\rm eff} > 700$ K. SIRTF, by contrast, is sensitive to the far more numerous objects at lower temperatures.

Fig. 2.— Synthetic spectra of isolated superplanets with 10 Jupiter mass at 10 pc from the Sun and a variety of ages. The detection limits of 2MASS (J, H and K_s) and our proosed SIRTF/IRAC survey are overplotted.

Fig. 3.— Smoothed model spectra for clear (blue) and cloudy (red) 30 $M_{\rm Jup}$ brown dwarf on arbitrary, linear flux scale (Marley 2000). Note the prominence of the $4 - 5\mu$m flux window. The clear model, like Gliese 229 B, is blue in J–K ($J - K \sim -1.6^m$) whereas the cloudy model is red ($J - K \sim 0.6^m$). Details of spectra are dependent upon uncertain cloud properties and will provide substantial information on cloud physics in a substellar mass atmosphere (Ackerman & Marley 2000). Flux units are in erg $\rm cm^{-2}$ $\rm sec^{-1}$ $\rm Hz^{-1}$.

Fig. 4.— In color-color space, brown dwarfs are clearly distinct from stellar sources (top panel) and extragalactic sources (lower panel).

Fig. 5.— Cumulative number of low-mass brown dwarfs and isolated superplanets that could be identified in our survey. Solid histrogram corresponds to a power-law IMF with $\alpha = 1.5$; Dotted histrogram corresponds to a power-law IMF with $\alpha = 1.0$.

Fig. 6.— Cumulative number of brown dwarfs and isolated superplanets identified at 5σ limiting flux in channel 2 as a function of effective temperature. We compare with a hypothetical survey that would require the same amount of SIRTF time (600 hours), covering more area but with less sensitivity. We have used an IMF with α=1.5.

This figure "fig1.jpg" is available in "jpg" format from:

http://arXiv.org/ps/astro-ph/0010315v1


4
2
0
−2
−4
log10 Fν [μJy]
1
10
λ[μm]
10 Mjup
0.1 Gyr (871 K)
0.5 Gyr (535 K)
1.0 Gyr (437 K)
5.0 Gyr (275 K)
2MASS
SIRTF


1.5×10^{-7}
10^{-7}
5×10^{-8}
0
Flux
1
2
3
4
5
λ (μm)
M ~
T$_{eff}$ =


$A_V = 20^m$
pop III brown dwarfs
flared disk
flat disk
5 M_{Jup}
10 M_{Jup}
20 M_{Jup}
30 M_{Jup}
Vega
pop I brown dwarfs
stars
0.1 Gyr
0.5 Gyr
1 Gyr
5 Gyr
1.0
0.5
0.0
-0.5
-1.0
-1.5
0
1
2
3
log F_ν(8.0μm)/F_ν(4.5μm)
log F_ν(4.5μm)/F_ν(3.6μm)


$A_V = 20^m$
pop III brown dwarfs
30 Dor
L* galaxies
z = 0.1 to 5
5 M_{Jup}
10 M_{Jup}
20 M_{Jup}
30 M_{Jup}
pop I brown dwarfs
0.1 Gyr
0.5 Gyr
1 Gyr
5 Gyr
1.0
0.5
0.0
-0.5
-1.0
-1.5
0
1
2
3
log F_ν(8.0μm)/F_ν(4.5μm)
log F_ν(4.5μm)/F_ν(3.6μm)


Isolated
Planets
α = 1.5
D-burning
mass limit
α = 1.0
Brown
Dwarfs
Log (Cumulative number per deg²)
Mass (Jovian masses)
0
0.5
1
1.5
5
10
15
20
25


18 deg^2, F$_2$=3.9μJy (5σ)
64 deg^2, F$_2$=107μJy (5σ)
Log (Cumulative number of objects detected)
Effective Temperature (K)
0
1
2
3
0
200
400
600
800
1000
1200